

Commonly used or illustrative parameters

Generic parameters	Value
Radioactive half life [1]	59.4 Days
Origin [1]	Decay of ¹²⁴ Xe
Principal decay mode [1]	Electron capture
Specific activity [2]	6.3 x 10 ¹⁴ Bq/g
Freshwater Kd [3]	4.4 x 10 ³ L kg ⁻¹
Marine Kd [4]	2.0 x 10 ² L kg ⁻¹

Parameters useful for human assessments	Value
CR Pasture grass [3]	3.7 x 10 ⁻³
CR Freshwater fish [3]	6.5 x 10 ² L kg ⁻¹
CR Marine fish [4]	9.0 x 10 ⁰ L kg ⁻¹
F _f Cow meat [3]	6.7 x 10 ⁻³ d kg ⁻¹
F _m Cow milk [3]	5.4 x 10 ⁻³ d L ⁻¹
Human fractional absorption (f1) [5]	1.0
Inhalation dose coefficient [6]	3.8 x 10 ⁻¹⁰ Sv Bq ⁻¹
Ingestion Dose Coefficient [6]	1.5 x 10 ⁻⁸ Sv Bq ⁻¹
Biological half life for Human (adult) [7]	a: 12 days (80 days in thyroid)
Biological half life for Cow milk [8]	a: 1.0 day (100%)
EU Food intervention limit- Dairy [9]	500 Bq L ⁻¹ or Bq kg ⁻¹
EU Food intervention limit- Baby food [9]	150 Bq L ⁻¹ or Bq kg ⁻¹
EU Food intervention limit- Liquid [9]	500 Bq L ⁻¹ or Bq kg ⁻¹
EU Food intervention limit- Other food [9]	2000 Bq L ⁻¹ or Bq kg ⁻¹
EU Food intervention limit- Minor food [9]	20000 Bq L ⁻¹ or Bq kg ⁻¹

Commonly used or illustrative parameters

Parameters useful for wildlife assessments	Value
Terrestrial EMCL — Soil [10]	$2.11 \times 10^5 \text{ Bq kg}^{-1}$
Freshwater EMCL — Water [10]	$1.19 \times 10^0 \text{ Bq L}^{-1}$
Freshwater EMCL — Sediment [10]	$5.56 \times 10^4 \text{ Bq kg}^{-1}$
Marine EMCL — Water [10]	$6.99 \times 10^0 \text{ Bq L}^{-1}$
Marine EMCL — Sediment [10]	$1.69 \times 10^2 \text{ Bq kg}^{-1}$
CR Terrestrial mammal (rat) [10]	4.0×10^{-1}
CR Freshwater fish [10]	3.2×10^2
CR Freshwater mollusc [10]	8.3×10^1
CR Marine fish [10]	9.0×10^0
CR Marine mollusc [10]	8.8×10^3
Internal DCC Terrestrial mammal (rat) on soil [10]	$3.3 \times 10^{-5} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ whole organism
External DCC Terrestrial mammal (rat) in soil [10]	$4.7 \times 10^{-6} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ soil
External DCC Terrestrial mammal (rat) on soil [10]	$1.9 \times 10^{-6} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ soil
Internal DCC Marine fish (benthic) [10]	$3.3 \times 10^{-5} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ whole organism
External DCC Marine fish (benthic) in water [10]	$1.4 \times 10^{-5} \mu\text{Gy h}^{-1}/\text{Bq L}^{-1}$ water
External DCC Marine fish (benthic) at sediment interface [10]	$7.0 \times 10^{-6} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ sediment
Internal DCC Freshwater fish (pelagic) [10]	$3.60 \times 10^{-5} \mu\text{Gy h}^{-1}/\text{Bq kg}^{-1}$ whole organism
External DCC Freshwater fish in water [10]	$1.10 \times 10^{-5} \mu\text{Gy h}^{-1}/\text{Bq L}^{-1}$ water

All terms used in these tables are described and discussed in underlying documents accessed via the hyperlinks provided

Sources of data [reference list](#)

Data compiled: September 2012

Data updated : April 2015

www.radioecology-exchange.org